

Smith Normal Form and Acyclic Matrices

In-Jae Kim and Bryan L. Shader

Department of Mathematics

University of Wyoming

Laramie, WY 82071, USA

February 2, 2008

Abstract

An approach, based on the Smith Normal Form, is introduced to study the spectra of symmetric matrices with a given graph. The approach serves well to explain how the path cover number (resp. diameter of a tree T) is related to the maximum multiplicity occurring for an eigenvalue of a symmetric matrix whose graph is T (resp. the minimum number $q(T)$ of distinct eigenvalues over the symmetric matrices whose graphs are T). The approach is also applied to a more general class of connected graphs G , not necessarily trees, in order to establish a lower bound on $q(G)$.

Keywords: Smith Normal Form, Acyclic matrix, spectra

MSC: 05C50

1 Introduction

This paper concerns the relationship between the algebraic and geometric properties of a symmetric matrix and the combinatorial arrangement of its nonzero entries (i.e. its graph). We begin by establishing some basic graph theoretic notation and terminology that follows that in [5].

A *graph* G consists of a vertex set $V(G)$ and an *edge* set $E(G)$, where an edge is an unordered pair of distinct vertices of G . We use uv to denote the edge joining vertices u and v . If uv is an edge, then we say that u and v are *adjacent*, and that v is a *neighbor* of u . A vertex is *incident* with an edge if it is one of the two vertices of the edge. The *degree* of a vertex is the number of edges incident to the vertex. A *subgraph* of a graph G is a graph H such that $V(H) \subseteq V(G)$, and $E(H) \subseteq E(G)$. A subgraph H is an *induced subgraph* if two vertices of $V(H)$ are adjacent in H if and only if they are adjacent in G . If $U \subseteq V(G)$, then $G \setminus U$ denotes the induced subgraph of G whose vertex set is $V(G) \setminus U$.

A *path* P of G is a sequence $v_1, v_2, \dots, v_n$ of distinct vertices such that consecutive vertices are adjacent, and is denoted by $v_1 - v_2 - \dots - v_n$. We say that P *covers* the vertices $v_1, \dots, v_n$, and that v_j is *covered by* P for $j = 1, \dots, n$. The *length* of P is the number of edges in P . If each vertex of G belongs to at most one of the paths $P_1, \dots, P_k$, then $P_1, \dots, P_k$ are *disjoint paths*.

If there is a path between each pair of vertices of G , then G is *connected*; otherwise G is *disconnected*. A *cycle* is a connected graph where every vertex has exactly two neighbors. The *length* of a cycle is the number of edges in the cycle. A *cycle in a graph* G is a subgraph

of G that is a cycle. An *acyclic graph* is a graph with no cycles. A connected acyclic graph is called a *tree*, and an acyclic graph is called a *forest*.

Let G be a connected graph. The *distance* between two vertices u and v of G is the minimum number of edges in a path from u to v . The *diameter* of a connected graph G is the maximum of the distances over pairs of vertices of G , and is denoted by $d(G)$. If G is a tree, $d(G)$ is the longest length of a path in G . The *path cover number* of G is the minimum number of disjoint paths needed to cover all of the vertices of G , and is denoted by $p(G)$.

As is customary, we use graphs to model the combinatorial structure of a matrix. Let $A = [a_{ij}]$ be an n by n symmetric matrix. The *graph* $G(A)$ of A consists of the vertices $1, 2, \dots, n$, and the edges ij for which $i \neq j$ and $a_{ij} \neq 0$. Note that $G(A)$ does not depend on the diagonal entries of A . An n by n symmetric matrix A is called an *acyclic* matrix if $G(A)$ is a tree (see [4]).

For a given graph G on n vertices, define $S(G)$ to be the set of all n by n real, symmetric matrices with graph G , i.e.

$$S(G) = \{A_{n \times n} \mid A \text{ is real, symmetric and } G(A) = G\}.$$

For the remainder of this section, matrices are real. Let σ be a multi-list of n real numbers. If there exists an n by n symmetric matrix A whose spectrum is σ , then we say that σ is *realized* by A , or A *realizes* σ . The *spectrum* of $S(G)$ for a graph G is the set of all spectra realized by some matrix in $S(G)$. For a given graph G , one can ask to characterize the spectrum of $S(G)$. This characterization problem is known as the Inverse Eigenvalue Problem for graphs, or IEP-G for short. If G is a tree, then we use IEP-T instead of IEP-G.

The IEP-G seems quite difficult. A first step toward resolving the IEP-G for a given graph G is to analyze the possible multiplicities of the eigenvalues in the spectra of matrices in $S(G)$. If the distinct eigenvalues of A are $\lambda_1 < \lambda_2 < \dots < \lambda_q$ and their corresponding multiplicities $m_1, \dots, m_q$, then $\langle m_1, m_2, \dots, m_q \rangle$ is the *ordered multiplicity list* of the eigenvalues of A .

The interplay between the spectral properties of acyclic matrices and the combinatorial properties of trees has been a fruitful area of research for the past 40 years, and some significant, intriguing and recent progress has been made on the IEP-T (see [4, 8, 9, 11, 12, 13, 14, 15]). The significant graphical parameters of trees T considered for the IEP-T are $d(T)$ and $p(T)$. For instance, A. Leal Duarte and C.R. Johnson showed in [8, 14] that the minimum number $q(T)$ of distinct eigenvalues over the matrices in $S(T)$ satisfies $q(T) \geq d(T) + 1$, and the maximum multiplicity $M(T)$ occurring for an eigenvalue of a matrix in $S(T)$ is $p(T)$.

In this paper, we introduce an approach based on the Smith Normal Form to study the spectra of matrices in $S(G)$. In Section 2 we relate the multiplicities of the eigenvalues of an n by n symmetric matrix A to the Smith Normal Form of $xI - A$ where I is the identity matrix. In Section 3 we provide a description of the determinant of an n by n matrix in terms of digraphs, and use the description to show that an eigenvalue of multiplicity $k + 1$ or more of an acyclic matrix $A \in S(T)$ for a tree T must be an eigenvalue of each principal submatrix of A whose graph is obtained from T by deleting k disjoint paths. In addition, as an application, we give an example showing that the IEP-T is not equivalent to determining the ordered multiplicity lists of the eigenvalues of matrices in $S(T)$. In Section 4, we show that the tight upper bound $p(T)$ on $M(T)$ is a direct consequence of the Smith Normal Form approach, and we describe a systematic way to compute $p(T)$ for a tree T . In Section 5, the bound $q(T) \geq d(T) + 1$ is easily derived, and we show that $q(W) \geq \frac{9d(W)}{8} + \frac{1}{2}$ for an infinite family of trees. In Section 6, we give a lower bound on $q(G)$ for a class of connected graphs G .

2 SNF and Multiplicities of Eigenvalues

Throughout the remainder of this paper, we let I denote an identity matrix of an appropriate order. In this section we give some useful results on the Smith Normal Form of matrices over the real polynomial ring, $\mathbb{R}[x]$. We refer the reader to [2, 6] for the basic facts. In particular, we relate the multiplicities of the eigenvalues of a real, symmetric matrix A to the Smith Normal Form of $xI - A$.

For $p(x), q(x) \in \mathbb{R}[x]$ we write $p(x) \mid q(x)$ if $p(x)$ divides $q(x)$, and $p(x) \nmid q(x)$ if $p(x)$ does not divide $q(x)$. We write $(x-a)^k \parallel q(x)$ provided $(x-a)^k \mid q(x)$ but $(x-a)^{k+1} \nmid q(x)$. We let $\mathbb{F}$ denote the field of rational functions over $\mathbb{R}$ (that is, $\mathbb{F}$ is the field of quotients of $\mathbb{R}[x]$), and let $(\mathbb{R}[x])^{m \times n}$ denote the set of all m by n matrices over $\mathbb{R}[x]$. Each $M \in (\mathbb{R}[x])^{m \times n}$ can be viewed as a matrix over $\mathbb{F}$, and the *rank* of M is defined to be the rank of M over $\mathbb{F}$. Let GL_n be the set of all invertible matrices of order n over $\mathbb{R}[x]$, i.e.

$$\text{GL}_n = \{P \in (\mathbb{R}[x])^{n \times n} \mid P \text{ has an inverse in } (\mathbb{R}[x])^{n \times n}\}.$$

It is a well-known fact that

$$\text{GL}_n = \{P \in \mathbb{R}[x]^{n \times n} \mid \det P \in \mathbb{R} \setminus \{0\}\}. \quad (1)$$

The matrices M and N in $(\mathbb{R}[x])^{m \times n}$ are *equivalent over $\mathbb{R}[x]$* if there exist $P \in \text{GL}_m$ and $Q \in \text{GL}_n$ such that $N = PMQ$. Hence equivalent matrices have the same rank.

A k by k *minor* of M is the determinant of a k by k submatrix of M . The monic greatest common divisor of all k by k minors of M is the k th *determinantal divisor* of M and is denoted by $\Delta_k(M)$. Another basic fact is:

Proposition 1 *Let $M, N \in (\mathbb{R}[x])^{n \times n}$. If M and N are equivalent over $\mathbb{R}[x]$, then $\Delta_k(M) = \Delta_k(N)$ for each $k \in \{1, \dots, n\}$.*

The following fundamental theorem asserts that each square matrix over $\mathbb{R}[x]$ is equivalent to a diagonal matrix over $\mathbb{R}[x]$ of a special form (see [2, 6]).

Theorem 2 (Smith Normal Form) *Let $M \in (\mathbb{R}[x])^{n \times n}$ of rank r . Then there exist $P, Q \in \text{GL}_n$ and monic polynomials $e_i(x)$ ($i = 1, 2, \dots, r$) such that $PMQ = D \oplus O$, where O is the zero matrix of order $n-r$, $D = \text{diag}(e_1(x), \dots, e_r(x))$, and $e_i(x) \mid e_{i+1}(x)$ for $i = 1, \dots, r-1$.*

Moreover, $\Delta_k(M) = \prod_{j=1}^k e_j(x)$ and $e_k(x) = \frac{\Delta_k(M)}{\Delta_{k-1}(M)}$ for each $k \in \{1, \dots, r\}$.

In the above theorem, $D \oplus O$ is called the *Smith Normal Form (SNF)* of M , and $e_i(x)$ is called the i th *invariant factor* of M .

Now assume that A is an n by n real matrix, and let S be the SNF of $xI - A$. The characteristic polynomial of A , denoted by $p_A(x)$, is $\det(xI - A) = \Delta_n(xI - A)$. Since $\det(xI - A)$ is nonzero, the rank of $xI - A$ is n . Thus, S is the full rank matrix of the form $\text{diag}(e_1(x), \dots, e_n(x))$. Since S and $xI - A$ are unimodularly equivalent, by Proposition 1,

$$p_A(x) = \prod_{j=1}^n e_j(x) = \Delta_n(S). \quad (2)$$

Further assume that A is symmetric. Then the spectrum of a symmetric matrix A , and the invariant factors of $xI - A$ are closely related. Let $P, Q \in \text{GL}_n$ such that $P(xI - A)Q = S$. Since A is symmetric, there exists an orthogonal matrix U of order n so that $U^T A U$ is $D = \text{diag}(\lambda_1, \dots, \lambda_n)$. The diagonal matrix D is called a *diagonalization* of A . Thus, $S = PU(xI - D)U^T Q$. Moreover, since (1) implies that $PU, U^T Q \in \text{GL}_n$, the SNF of $xI - D$ is also S . This along with Proposition 1 imply that $\Delta_i(xI - A) = \Delta_i(xI - D) = \Delta_i(S)$ for all i .

Henceforth $\Delta_i(x)$ denotes $\Delta_i(xI - A)$. If λ is an eigenvalue of A , then $m_A(\lambda)$ denotes the algebraic multiplicity of λ . Let $p(x) \in \mathbb{R}[x]$, and $a \in \mathbb{R}$. If $(x - a) \parallel p(x)$, then $x - a$ is a linear factor of $p(x)$. By considering $xI - D$, we have the following result on the factors of $\Delta_i(x)$ and $e_i(x)$.

Theorem 3 *Let A be an n by n symmetric matrix whose distinct eigenvalues are $\lambda_1, \lambda_2, \dots, \lambda_q$, and let $S = \text{diag}(e_1(x), \dots, e_n(x))$ be the SNF of $xI - A$. Then the following hold:*

- (a) *if $k \leq n - m_A(\lambda_j)$, then $(x - \lambda_j) \nmid \Delta_k(x)$ and $(x - \lambda_j) \nmid e_k(x)$,*
- (b) *if $k > n - m_A(\lambda_j)$, then $(x - \lambda_j)^{k-n+m_A(\lambda_j)} \parallel \Delta_k(x)$ and $x - \lambda_j$ is a linear factor of $e_k(x)$,*
- (c) $e_{n-k}(x) = \prod_{j: m_A(\lambda_j) > k} (x - \lambda_j)$

Proof. Fix j , and let $\lambda = \lambda_j$, and $m = m_A(\lambda_j)$. Let D be a diagonalization of A . Without loss of generality, we may assume

$$xI - D = \begin{bmatrix} x - \mu_1 & & \\ & \ddots & \\ & & x - \mu_{n-m} \end{bmatrix} \oplus \begin{bmatrix} x - \lambda & & \\ & \ddots & \\ & & x - \lambda \end{bmatrix}_{m \times m},$$

where $\mu_i \neq \lambda$ for each $i = 1, \dots, n - m$.

(a) Suppose $k \leq n - m$. The determinant of $\text{diag}(x - \mu_1, \dots, x - \mu_k)$ is not divisible by $x - \lambda$. Thus, $(x - \lambda) \nmid \Delta_k(x)$. By Theorem 2, $e_k(x) \mid \Delta_k(x)$. Hence, $(x - \lambda) \nmid e_k(x)$.

(b) Suppose $k > n - m$, and let M be a k by k submatrix of $xI - D$. If M is not a principal submatrix of $xI - D$, then M has a zero row and hence $\det M = 0$. Otherwise, at least $k - (n - m)$ diagonal entries of M are $x - \lambda$. Thus, $(x - \lambda)^{k-(n-m)} \mid \Delta_k(x)$. Note that $xI - D$ has a k by k minor equal to $\det \text{diag}(x - \mu_1, \dots, x - \mu_{n-m}, x - \lambda, \dots, x - \lambda)$. Thus $(x - \lambda)^{k-(n-m)+1} \nmid \Delta_k(x)$. Hence

$$(x - \lambda)^{k-(n-m)} \parallel \Delta_k(x). \quad (3)$$

By Theorem 2, $e_k(x) = \frac{\Delta_k(x)}{\Delta_{k-1}(x)}$. By (3), $\Delta_k(x)$ has exactly $k - n + m$ factors equal to $x - \lambda$ and $\Delta_{k-1}(x)$ has exactly $(k - n + m) - 1$ factors equal to $x - \lambda$. Thus $(x - \lambda) \parallel e_k(x)$, and (b) holds.

(c) By (2), $e_{n-k}(x)$ is a product of linear factors from $\{x - \lambda_1, \dots, x - \lambda_q\}$, and by (a) and (b), the factors are distinct, and $x - \lambda_j$ is a factor of $e_{n-k}(x)$ if and only if $m_A(\lambda_j) > k$. Thus (c) holds. $\blacksquare$

Useful, immediate consequences of Theorem 3 are the following:

Corollary 4 *Let A be an n by n symmetric matrix, and $S = \text{diag}(e_1(x), \dots, e_n(x))$ be the SNF of $xI - A$. Suppose that λ is an eigenvalue of A . Then*

- (a) $m_A(\lambda) \geq k$ if and only if $(x - \lambda) \mid e_{n-k+1}$
- (b) $m_A(\lambda) = k$ if and only if $(x - \lambda) \mid e_{n-k+1}(x)$ but $(x - \lambda) \nmid e_{n-k}(x)$.

Corollary 4 implies

$$\deg(e_{n-k}(x)) \text{ is the number of eigenvalues of } A \text{ with multiplicity } k + 1 \text{ or more.} \quad (4)$$

Taking $k = 0$ in (c) of Theorem 3, we see that $e_n = (x - \lambda_1)(x - \lambda_2) \cdots (x - \lambda_q)$, which is known to be the minimal polynomial of A . Thus, $\deg(e_n)$ equals the number $q(A)$ of distinct eigenvalues of A , and by Theorem 2, we have the following:

Corollary 5 *Let A be an n by n symmetric matrix, and $S = \text{diag}(e_1(x), \dots, e_n(x))$ be the SNF of $xI - A$. Then $e_n(x)$ is the minimal polynomial of A , and*

$$q(A) = n - \deg(\Delta_{n-1}(x)).$$

Corollary 5 allows one to obtain a lower bound on $q(A)$ from an upper bound on $\deg(\Delta_{n-1}(x))$.

3 Eigenvalues of Principal Submatrices of an acyclic matrix

In this section we associate a digraph on n vertices with an n by n matrix M over $\mathbb{R}[x]$, and describe $\det M$ in terms of the structure of the digraph associated with M . We use this description to show that an eigenvalue of a symmetric acyclic matrix A with multiplicity $k + 1$ or more is an eigenvalue of each principal submatrix of A whose indices correspond to the vertices not covered by a set of k disjoint paths. As an application, we provide an example showing that the IEP-T is not equivalent to determining the ordered multiplicity lists of the eigenvalues of matrices in $S(T)$ (see also [1]).

First, we give some necessary definitions. Let T be a tree, Q an induced subgraph of T , and M a symmetric matrix over $\mathbb{R}[x]$ with $G(M) = T$. Then $M[Q]$ denotes the principal submatrix of M whose rows and columns correspond to the vertices of Q , and $T \setminus Q$ denotes the induced subgraph of T obtained by deleting all vertices of Q . If $Q_1, \dots, Q_k$ are induced subgraphs of T , $Q_1 \cup \dots \cup Q_k$ denotes the induced subgraph of T whose vertex set is the union of the vertex sets of $Q_1, \dots, Q_k$.

The *end vertices* of the path $v_1 - v_2 - \dots - v_\ell$ are v_1 and v_ℓ . If $P_1 = v_1 - \dots - v_s$ and $P_2 = v_s - \dots - v_t$ are paths whose only common vertex is v_s , then $P_1 P_2$ denotes the path $v_1 - \dots - v_s - \dots - v_t$, obtained by concatenating P_1 and P_2 .

Let $M = [m_{ij}]$ be an n by n matrix over $\mathbb{R}[x]$. The *digraph* $D(M)$ of M consists of the vertices $1, 2, \dots, n$, and arcs (i, j) from vertex i to vertex j if and only if $m_{ij} \neq 0$. An arc from a vertex to itself is called a *loop*. A *subdigraph* H of $D(M)$ is a digraph such that the vertex set of H is a subset of $\{1, 2, \dots, n\}$ and the arc set of H is a subset of the arc set of $D(M)$. The *underlying graph* of the digraph $D(M)$ is the graph obtained by treating each arc (i, j) ($i \neq j$) of $D(M)$ as the edge ij , and ignoring the loops.

A *directed walk* in $D(M)$ is a sequence of vertices $(v_1, v_2, \dots, v_\ell)$, such that (v_i, v_{i+1}) is an arc for each $i = 1, \dots, \ell - 1$, and v_1 is the *initial* vertex and v_ℓ is the *terminal* vertex of the directed walk. The directed walk $(v_1, v_2, \dots, v_\ell)$ *covers* the vertices $v_1, v_2, \dots, v_\ell$, and has *length* $\ell - 1$. If W_1 is the directed walk $(v_1, \dots, v_s)$, and W_2 is the directed walk $(v_s, \dots, v_t)$, then (W_1, W_2) is the directed walk $(v_1, \dots, v_s, \dots, v_t)$. If no vertex of a directed walk is repeated, then the directed walk is a *directed path*.

If the underlying graph of $D(M)$ is a tree T , then there is at most one directed path from vertex i to vertex j . If there exists one, $P_{i \rightarrow j}$ denotes the unique directed path from vertex i to vertex j in $D(M)$. Let P_{i-j} denote the unique path connecting i and j in the underlying graph T of $D(M)$.

If the initial vertex is equal to the terminal vertex in a directed walk, then the directed walk is *closed*. A *directed cycle* is a closed directed walk with no repeated vertices other than the initial and terminal vertices. If a directed cycle has length r , the directed cycle is a *directed r -cycle*. If each vertex is incident to at most one of the directed cycles $\gamma_1, \dots, \gamma_k$ in D , then $\gamma_1, \dots, \gamma_k$ are *disjoint*.

The *weight of the arc* (i, j) of $D(M)$ is m_{ij} . The *weight of a directed walk* β of $D(M)$ is the product of the weights of its arcs, and is denoted by $\text{wt}(\beta)$. The *signed weight of a directed cycle* γ , denoted by $\text{swt}(\gamma)$, is $(-1)^{\ell-1} \text{wt}(\gamma)$ where ℓ is the length of γ . Let

$\alpha = \{\gamma_1, \dots, \gamma_t\}$ where $\gamma_1, \dots, \gamma_t$ are disjoint directed cycles in $D(M)$ covering all of the n vertices. We define the *signed weight* of α to be the product of the signed weights of $\gamma_1, \dots, \gamma_t$. Assume that Γ is the set of all such α 's described above. Then, by the definition of the determinant of a square matrix, it is known [3] (see p.291) that

$$\det M = \sum_{\alpha \in \Gamma} \text{swt}(\alpha). \quad (5)$$

Let M be an n by n matrix. If α and β are subsets of $\{1, \dots, n\}$, then we denote the submatrix of M obtained by removing (resp. retaining) rows indexed by α and columns indexed by β by $M(\alpha, \beta)$ (resp. $M[\alpha, \beta]$). When $\alpha = \beta$, we use $M(\alpha)$ and $M[\alpha]$, respectively. We use e_i to denote the i th column of the identity matrix I .

In the following theorem, we provide a relation between the determinant of a submatrix and the determinant of a principal submatrix of an acyclic matrix over $\mathbb{R}[x]$.

Theorem 6 *Let M be an n by n symmetric matrix over $\mathbb{R}[x]$, where the underlying graph of $D(M)$ is a tree T , and let $P_{i_1-j_1}, \dots, P_{i_k-j_k}$ be disjoint paths in T . Then*

$$\det M(\{j_1, \dots, j_k\}, \{i_1, \dots, i_k\}) = \pm \prod_{s=1}^k \text{wt}(P_{i_s \rightarrow j_s}) \cdot \det M[T \setminus (P_{i_1-j_1} \cup \dots \cup P_{i_k-j_k})].$$

Proof. Let M' be the matrix obtained from M by replacing the i_s th column of M by e_{j_s} for each $s = 1, \dots, k$. By the construction of M' , $M'(\{j_1, \dots, j_k\}, \{i_1, \dots, i_k\}) = M(\{j_1, \dots, j_k\}, \{i_1, \dots, i_k\})$. By Laplace expansion of the determinant along the columns $i_1, i_2, \dots, i_k$ of M' , we have

$$\begin{aligned} \det M' &= \pm \det M'(\{j_1, \dots, j_k\}, \{i_1, \dots, i_k\}) \\ &= \pm \det M(\{j_1, \dots, j_k\}, \{i_1, \dots, i_k\}). \end{aligned} \quad (6)$$

In terms of digraphs, $D(M')$ is obtained from $D(M)$ by deleting all incoming arcs to vertices $i_1, \dots, i_k$ and inserting the arcs $(j_1, i_1), \dots, (j_k, i_k)$, each with weight 1. Therefore, $D(M')$ has exactly one arc, namely (j_s, i_s) , ending at vertex i_s for each $s = 1, \dots, k$.

Set $U = \{(j_1, i_1), \dots, (j_k, i_k)\}$. We claim that each directed cycle in $D(M')$ has at most one arc in U . Suppose to the contrary that there exists a directed cycle γ with more than one arc in U . Without loss of generality, we may assume that $(j_1, i_1), \dots, (j_t, i_t)$ are arcs of γ in U , and $\gamma = ((j_1, i_1), P_{i_1 \rightarrow j_2}, \dots, (j_t, i_t), P_{i_t \rightarrow j_1})$ (see Figure 1).




Figure 1.

Since $P_{i_1 \rightarrow j_2}, \dots, P_{i_t \rightarrow j_1}$ are directed paths in the directed cycle γ , the underlying graphs of $P_{i_1 \rightarrow j_2}, \dots, P_{i_t \rightarrow j_1}$ are disjoint paths in T .

Next, we consider $P_{j_s \rightarrow i_s}$, the unique directed path from j_s to i_s in $D(M)$ for each $s = 1, \dots, t$. Then $\tau = (P_{j_1 \rightarrow i_1}, P_{i_1 \rightarrow j_2}, \dots, P_{j_t \rightarrow i_t}, P_{i_t \rightarrow j_1})$ is a directed closed walk in $D(M)$. Since the underlying graph of $D(M)$ is T , the multi-set of the arcs in τ is the disjoint union of the sets of the arcs of directed 2-cycles. Note that $P_{j_1 \rightarrow i_1}, \dots, P_{j_t \rightarrow i_t}$ are disjoint and that $P_{i_1 \rightarrow j_2}, \dots, P_{i_t \rightarrow j_1}$ are disjoint. Thus, the disjoint union of the sets of the edges of $P_{j_1 \rightarrow i_1}, \dots, P_{j_t \rightarrow i_t}$ is equal to the disjoint union of the sets of the edges of $P_{i_1 \rightarrow j_2}, \dots, P_{i_t \rightarrow j_1}$. Since the paths $P_{j_1 \rightarrow i_1}, \dots, P_{j_t \rightarrow i_t}$ in T are disjoint, there is no path from i_1 to j_2 , consisting of the arcs from $P_{j_1 \rightarrow i_1}, \dots, P_{j_t \rightarrow i_t}$. This contradicts that $P_{i_1 \rightarrow j_2}$ is a directed path from i_1 to j_2 . Therefore, each directed cycle in $D(M')$ has at most one arc in U .

This implies that each set of disjoint directed cycles of $D(M')$ that cover every vertex consists of the directed cycles $\beta_s = ((j_s, i_s), P_{i_s \rightarrow j_s})$ for all $s = 1, \dots, k$ along with disjoint directed cycles covering every vertex of $T \setminus (P_{i_1 \rightarrow j_1} \cup \dots \cup P_{i_k \rightarrow j_k})$. Therefore, (5) implies that

$$\det M' = \prod_{s=1}^k \text{swt}(\beta_s) \cdot \sum_{\alpha \in \Gamma(M'[T \setminus (P_{i_1 \rightarrow j_1} \cup \dots \cup P_{i_k \rightarrow j_k})])} \text{swt}(\alpha).$$

Since $M'[T \setminus (P_{i_1 \rightarrow j_1} \cup \dots \cup P_{i_k \rightarrow j_k})] = M[T \setminus (P_{i_1 \rightarrow j_1} \cup \dots \cup P_{i_k \rightarrow j_k})]$, (5) implies

$$\det M' = \prod_{s=1}^k \text{swt}(\beta_s) \cdot \det M[T \setminus (P_{i_1 \rightarrow j_1} \cup \dots \cup P_{i_k \rightarrow j_k})].$$

Note that $\text{swt}(\beta_s) = \pm \text{wt}(P_{i_s \rightarrow j_s})$ for each $s = 1, \dots, k$. Hence, by (6),

$$\begin{aligned} \det M(\{j_1, \dots, j_k\}, \{i_1, \dots, i_k\}) &= \pm \det M' \\ &= \pm \prod_{s=1}^k \text{wt}(P_{i_s \rightarrow j_s}) \cdot \det M[T \setminus (P_{i_1 \rightarrow j_1} \cup \dots \cup P_{i_k \rightarrow j_k})]. \end{aligned}$$

■

Corollary 7 Let $A \in S(T)$, where T is a tree on n vertices, and let $S = \text{diag}(e_1(x), \dots, e_n(x))$ be the SNF of $xI - A$. If $P_{i_1 \rightarrow j_1}, \dots, P_{i_k \rightarrow j_k}$ are disjoint paths in T covering $n - t$ vertices of T , then

$$\Delta_{n-k}(x) | \det((xI - A)[T \setminus (P_{i_1 \rightarrow j_1} \cup \dots \cup P_{i_k \rightarrow j_k})]) \text{ and } \deg(\Delta_{n-k}(x)) \leq t.$$

Furthermore, if λ is an eigenvalue of A with $m_A(\lambda) \geq k + 1$, then λ is an eigenvalue of $A[T \setminus (P_{i_1 \rightarrow j_1} \cup \dots \cup P_{i_k \rightarrow j_k})]$ with multiplicity $m_A(\lambda) - k$ or more.

Proof. Let $P_{i_s \rightarrow j_s}$ be the directed path from i_s to j_s in $D(xI - A)$ whose underlying graph is $P_{i_s \rightarrow j_s}$ for each $s = 1, \dots, k$. By Theorem 6,

$$\det[(xI - A)(\{j_1, \dots, j_k\}, \{i_1, \dots, i_k\})] = \pm \prod_{s=1}^k \text{wt}(P_{i_s \rightarrow j_s}) \cdot \det((xI - A)[T \setminus (P_{i_1 \rightarrow j_1} \cup \dots \cup P_{i_k \rightarrow j_k})]).$$

Note that $\text{wt}(P_{i_s \rightarrow j_s})$ is a nonzero constant for each $s = 1, \dots, k$. Hence,

$$\det[(xI - A)(\{j_1, \dots, j_k\}, \{i_1, \dots, i_k\})] = c \cdot \det((xI - A)[T \setminus (P_{i_1 \rightarrow j_1} \cup \dots \cup P_{i_k \rightarrow j_k})])$$

for some nonzero constant c .

Since $(xI - A)(\{j_1, \dots, j_k\}, \{i_1, \dots, i_k\})$ is an $n - k$ by $n - k$ submatrix of $xI - A$,

$$\Delta_{n-k}(x) | \det((xI - A)[T \setminus (P_{i_1 \rightarrow j_1} \cup \dots \cup P_{i_k \rightarrow j_k})]). \quad (7)$$

By Theorem 3, if $m_A(\lambda) \geq k+1$, then the multiplicity of $(x-\lambda)$ as a factor of $\Delta_{n-k}(x)$ is $m_A(\lambda)-k$. Therefore, by (7), $(x-\lambda)^{m_A(\lambda)-k} \mid \det((xI-A)[T \setminus (P_{i_1-j_1} \cup \dots \cup P_{i_k-j_k})])$. This implies that λ is an eigenvalue of $A[T \setminus (P_{i_1-j_1} \cup \dots \cup P_{i_k-j_k})]$ with multiplicity $m_A(\lambda)-k$ or more.

Note that the degree of $\det((xI-A)[T \setminus (P_{i_1-j_1} \cup \dots \cup P_{i_k-j_k})])$ is t . Thus, by (7),

$$\deg(\Delta_{n-k}(x)) \leq t.$$

■

If $\deg(\Delta_{n-k}(x)) \leq t$, then Theorem 2 implies $\deg(e_{n-k}(x)) \leq t$. Hence, Corollary 4 implies the following.

Corollary 8 *Let $A \in S(T)$, where T is a tree on n vertices. If k disjoint paths of T cover $n-t$ vertices of T , then there are at most t eigenvalues of A with multiplicity $k+1$ or more.*

It was conjectured in [11] that the IEP-T for a tree T is equivalent to determining the ordered multiplicity lists of the eigenvalues of matrices in $S(T)$, i.e. each multi-list of real numbers having an ordered multiplicity list of the eigenvalues of a matrix in $S(T)$ is the spectrum of a matrix in $S(T)$. Indeed, [11] showed that for some classes of trees, these two problems are equivalent. A counterexample to the conjecture was given in [1]. We give a counterexample on fewer vertices, and a simple argument that shows this is a counterexample.

Consider the tree T illustrated in Figure 2. We will show that an ordered multiplicity list of the eigenvalues of a matrix in $S(T)$ requires the eigenvalues having the ordered multiplicity list to satisfy a certain algebraic condition.




Figure 2.

Example 9 *It can be verified that the eigenvalues of the following matrix A in $S(T)$ are $-\sqrt{5}$, $-\sqrt{2}$, 0 , $\sqrt{2}$, $\sqrt{5}$, and the ordered multiplicity list of the eigenvalues of A is $\langle 1, 2, 4, 2, 1 \rangle$:*

$$\left[\begin{array}{ccc|cccccc} 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 1 \\ \hline 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

Suppose that

$$\sigma = (\lambda_1, \lambda_2, \lambda_2, \lambda_3, \lambda_3, \lambda_3, \lambda_3, \lambda_4, \lambda_4, \lambda_5)$$

is realized by a matrix B in $S(T)$ as its spectrum where $\lambda_1 < \lambda_2 < \lambda_3 < \lambda_4 < \lambda_5$.

The disjoint paths $4-1-5$, $7-2-8$; and $10-3-9$ cover all the vertices except vertex 6 (see Figure 3).




Figure 3.

Since $m_B(\lambda_3) = 4$, Corollary 7 implies that λ_3 is the eigenvalue of $B[6]$, i.e. $B[6] = \lambda_3$.

Next, since the three disjoint paths $4-1-6-2-7$; $10-3-9$; and 8 cover all the vertices except vertex 5 (see Figure 4), Corollary 7 implies that λ_3 is the eigenvalue of $B[5]$, i.e. $B[5] = \lambda_3$.




Figure 4.

Similarly, λ_3 is the eigenvalue of $B[i]$, i.e. $B[i] = \lambda_3$ for each $i = 4, 7, 8, 9, 10$.

The three disjoint paths $7-2-6-3-10$; 8 and 9 cover seven vertices (see Figure 5).




Figure 5.

Thus, by Corollary 7, λ_3 is an eigenvalue of $B[\{4, 1, 5\}]$. Similarly, λ_3 is an eigenvalue of $B[\{10, 3, 9\}]$ and $B[\{7, 2, 8\}]$.

Finally, we show that both eigenvalues of B with multiplicity 2 are eigenvalues of $B[\{4, 1, 5\}]$, $B[\{7, 2, 8\}]$ and $B[\{10, 3, 9\}]$. We consider only the case for $B[\{4, 1, 5\}]$. Since the single path $7-2-6-3-10$ covers five vertices, and $m_B(\lambda_2) = m_B(\lambda_4) = 2$, Corollary 7 implies that λ_2 and λ_4 are eigenvalues of $B[\{4, 1, 5, 8, 9\}]$. However, $B[\{4, 1, 5, 8, 9\}] = B[\{4, 1, 5\}] \oplus B[8] \oplus B[9]$. Since $B[8] = B[9] = \lambda_3$, λ_2 and λ_4 are eigenvalues of $B[\{4, 1, 5\}]$. The other cases can be shown by choosing the paths $4-1-6-2-7$ and $4-1-6-3-10$, respectively.

So far, we have shown that $\sigma(B[6]) = \lambda_3$, and $\sigma(B[\{4, 1, 5\}]) = \sigma(B[\{7, 2, 8\}]) = \sigma(B[\{10, 3, 9\}]) = (\lambda_2, \lambda_3, \lambda_4)$. Now, we consider the trace of B . The trace of B is equal to the sum of the traces of $B[6]$, $B[\{4, 1, 5\}]$, $B[\{7, 2, 8\}]$ and $B[\{10, 3, 9\}]$. Since the trace

is equal to the sum of all the eigenvalues, we have

$$\lambda_1 + 2\lambda_2 + 4\lambda_3 + 2\lambda_4 + \lambda_5 = \lambda_3 + 3(\lambda_2 + \lambda_3 + \lambda_4)$$

and hence,

$$\lambda_1 + \lambda_5 = \lambda_2 + \lambda_4.$$

Therefore, if the ordered multiplicity list $\langle 1, 2, 4, 2, 1 \rangle$ is realized by a matrix B in $S(T)$, then $\sigma(B) = (\lambda_1, \lambda_2, \lambda_2, \lambda_3, \lambda_3, \lambda_3, \lambda_3, \lambda_4, \lambda_4, \lambda_5)$ must satisfy $\lambda_1 + \lambda_5 = \lambda_2 + \lambda_4$. For instance, $\sigma = (2, 3, 3, 5, 5, 5, 5, 7, 7, 10)$ with the ordered multiplicity list $\langle 1, 2, 4, 2, 1 \rangle$ cannot be realized by any matrix in $S(T)$.

4 Maximum Multiplicity and $p(T)$

Let T be a tree on n vertices, and let $M(G)$ to denote the maximum multiplicity occurring for an eigenvalue among the matrices in $S(T)$. Recall that the path cover number of T is the minimum number of disjoint paths that cover T .

Now let $A \in S(T)$. If k disjoint paths in T cover all the vertices of T , then Corollary 8 implies that no eigenvalue of A has multiplicity more than k . Since A is an arbitrary matrix in $S(T)$, we have the following corollary.

Corollary 10 *Let T be a tree. Then*

$$M(T) \leq p(T).$$

Furthermore, it was shown in [8] that the upper bound is tight.

Theorem 11 *Let T be a tree. Then*

$$M(T) = p(T).$$

For the remainder of this section, we describe a systematic way of computing $p(T)$ for a tree T . This method will be used repeatedly in the following sections. We first show that the existence of a specific path for a given tree (see also [15, Lemma 3.1]).

Proposition 12 *Let T be a tree on n vertices. Then there exists a path in T such that the end vertices of the path are pendant vertices of T , and at most one vertex of the path has degree 3 or more in T .*

Proof. The proof is by induction on n . The result is clear if $n \leq 2$. Assume $n \geq 3$ and proceed by induction.

If T has diameter 2, then any path of length 2 works. Assume that the diameter of T is at least 3, and let P be a path in T , $u-v-a_1-\dots-a_{k-1}-a_k$, whose length is the diameter of T . Since u is a pendant vertex of T , $T \setminus \{u\}$ is also a tree. By the inductive hypothesis, there exists a path P' of $T \setminus \{u\}$ satisfying the given condition.

If the path P' in $T \setminus \{u\}$ does not contain v , then P' is also a path in T satisfying the given condition.

If P' contains v as an end vertex, then the path $P_{u-v}P'$ is a path in T satisfying the given condition.

Otherwise P' contains v and v is not an end vertex of P' . Thus there exists a neighbor w of v in $T \setminus \{u\}$ such that $w \neq a_1$. Suppose that w is not a pendant vertex of T . Then there exists a neighbor y of w other than v . Since T is a tree $y \notin \{a_2, \dots, a_k\}$ and hence, $y-w-v-a_1-\dots-a_{k-1}-a_k$ is a path in T , whose length is longer than P , which is a contradiction. Therefore, w is a pendant vertex of T , and the path $u-v-w$ in T satisfies the given condition. ■

If T is a path, then a path in T satisfying the conditions in Proposition 12 is T itself. Otherwise, there exists a path P in T such that the end vertices of P are pendant vertices of T , and exactly one vertex of P has degree 3 or more in T . Next, we show that for such a path P , $p(T \setminus P) = p(T) - 1$.

Proposition 13 *Let T be a tree which is not a path. Suppose that P is a path in T such that P 's end vertices are pendant vertices of T and P has exactly one vertex v of degree 3 or more in T . Then*

$$p(T) = p(T \setminus P) + 1.$$

Proof. Note that each path cover of $T \setminus P$ can be extended to a path cover of T by including the path P . Hence $p(T) \leq p(T \setminus P) + 1$

Now we show $p(T) \geq p(T \setminus P) + 1$. Let p denote $p(T)$. and $C = \{P_i\}_{i=1}^p$ be a set of p disjoint paths in T covering all of the vertices of T .

If $P \in C$, then, since $C \setminus \{P\}$ covers all of the vertices of $T \setminus P$, $p(T \setminus P) \leq p - 1$.

Otherwise, $P \notin C$. Then two disjoint paths in C , say α and β , are needed to cover the vertices of P . Assume that β covers the vertex v . Then α covers only the vertices of P . Thus, $(C \setminus \{\alpha, \beta\}) \cup \{\beta \setminus P\}$ covers all the vertices of $T \setminus P$. This implies that $p(T \setminus P) \leq p - 1$; equivalently, $p \geq p(T \setminus P) + 1$. ■

By repeated use of Proposition 13, we can effectively compute the path cover number of the tree T in Figure 6.




Figure 6.

Example 14 *Since $P_1, 3-1-4$, and $P_2, 9-2-10$, satisfy the condition in Proposition 13 for T and $T \setminus P_1$, respectively, we have the following three disjoint paths covering all of the vertices of T .*




Figure 7.

Hence, $p(T) = 3$. This implies, by Theorem 11, that if a multi-list σ of 10 real numbers has an element with multiplicity greater than 3, σ cannot be realized by any matrix in $S(T)$.

5 Relationship between $d(T)$ and $q(T)$

Let T be a tree, and $A \in S(T)$. The number of distinct eigenvalues of A is denoted by $q(A)$ and, $q(T)$ denotes the minimum of $q(A)$ over all $A \in S(T)$, and $d(T)$ is the diameter of T . In this section, we study the relation between $d(T)$ and $q(T)$.

Let $\Delta_i(x)$ be the i th determinantal divisor of $xI - A$. By Corollary 5,

$$q(A) = n - \deg(\Delta_{n-1}(x)) \quad (8)$$

and by Corollary 7, if a path in T has ℓ vertices, then

$$\deg(\Delta_{n-1}(x)) \leq n - \ell. \quad (9)$$

Thus, by choosing a path of the longest length, (8) and (9) imply that

$$q(A) \geq d(T) + 1. \quad (10)$$

Since (10) holds for every matrix in $S(T)$, $q(T) \geq d(T) + 1$. Thus, the following known theorem (see [14]) follows easily from the Smith Normal Form approach.

Theorem 15 *Let T be a tree. Then*

$$q(T) \geq d(T) + 1.$$

Next, we provide a class of trees W for which $q(W)$ is much larger than $d(W) + 1$. The $(3, \ell)$ -whirl ($\ell \geq 1$), W , is the tree on $n = 6\ell + 4$ vertices with 6 pendant paths $\alpha_i, \beta_i, \gamma_i$, each with ℓ vertices, for $i = 1, 2$ as illustrated in Figure 8. The vertex v in Figure 8 is the *axis vertex* of W , and each of the pendant paths is a *leg* of W .




Figure 8.

Note that $d(W) = 2\ell + 2$.

Theorem 16 *Let W be the $(3, \ell)$ -whirl with $\ell \geq 2$. Then*

$$q(W) \geq \frac{9d(W)}{8} + \frac{1}{2} = d(W) + 1 + \frac{\ell - 1}{4} > d(W) + 1.$$

Proof. Applying Proposition 13 to the four paths illustrated in Figure 9, we conclude $p(W) = 4$.




Figure 9.

Let $A \in S(W)$, and let n_j be the number of the eigenvalues of A with multiplicity j . Then, since $p(W) = 4$, Theorem 11 implies $n_j = 0$ for $j > 4$. The three disjoint paths P_1, P_2 and P_3 in W in Figure 9 cover all of the vertices of W except v . Thus, by Corollary 8,

$$n_4 \leq 1. \quad (11)$$

Suppose that $\lambda \in \sigma(A)$ and $m_A(\lambda) = 4$. Consider the three disjoint paths P_1, P_2 and P_3 in W illustrated in Figure 10.




Figure 10.

Then, by Corollary 7, λ is an eigenvalue of $A[W \setminus (P_1 \cup P_2 \cup P_3)] = A[\alpha_1]$. Similarly, λ is an eigenvalue of $A[\alpha_2]$, $A[\beta_i]$ and $A[\gamma_i]$ for $i = 1, 2$. Since α_i, β_i and γ_i are paths, Theorem 11 implies that λ is a simple eigenvalue of $A[\alpha_i]$, $A[\beta_i]$ and $A[\gamma_i]$ for each $i = 1, 2$.

Next, suppose $\mu \in \sigma(A)$ and $m_A(\mu) = 3$. Let P_1 and P_2 be the disjoint paths of W illustrated in Figure 11.




Figure 11.

Then, by Corollary 7, μ is an eigenvalue of $A[W \setminus (P_1 \cup P_2)]$ which is, up to permutation similarity, equal to $A[\alpha_1] \oplus A[\gamma_2]$. Thus, by Theorem 11, μ is a simple eigenvalue of at least one of $A[\alpha_1]$, $A[\gamma_2]$. Similarly, μ is a simple eigenvalue of $A[\alpha_i]$ or $A[\beta_j]$, $A[\alpha_i]$ or $A[\gamma_j]$,

and $A[\beta_i]$ or $A[\gamma_j]$ for $i, j \in \{1, 2\}$. If μ is a simple eigenvalue of at most three matrices in

$$\{A[\alpha_1], A[\beta_1], A[\gamma_1], A[\alpha_2], A[\beta_2], A[\gamma_2]\}, \quad (12)$$

then there is a pair of $A[\delta_i], A[\tau_j]$ for $\delta, \tau \in \{\alpha, \beta, \gamma\}$ and $\delta \neq \tau$ such that neither $A[\delta_i]$ nor $A[\tau_j]$ has μ as an eigenvalue. Thus, μ is a simple eigenvalue of at least four matrices in (12).

Let $A' = A[\alpha_1] \oplus A[\alpha_2] \oplus A[\beta_1] \oplus A[\beta_2] \oplus A[\gamma_1] \oplus A[\gamma_2]$. We have shown (a) if $\lambda \in \sigma(A)$ and $m_A(\lambda) = 4$, then $\lambda \in \sigma(A')$ and $m_{A'}(\lambda) = 6$; and (b) if $\mu \in \sigma(A)$ and $m_A(\mu) = 3$, then $\mu \in \sigma(A')$ and $m_{A'}(\mu) \geq 4$. Since the number of eigenvalues of a square matrix cannot exceed its order, by (a) and (b),

$$4n_3 + 6n_4 \leq 6\ell.$$

Hence,

$$n_3 \leq \frac{3\ell - 3n_4}{2}. \quad (13)$$

By definition of n_j and the fact that no eigenvalue of A has multiplicity more than 4,

$$n = n_1 + 2n_2 + 3n_3 + 4n_4 = n_1 + 2(n_2 + n_3 + n_4) + n_3 + 2n_4. \quad (14)$$

It is shown in [10, Corollary 7] that the largest and smallest eigenvalues of A are simple and hence, $n_1 \geq 2$. Since $n_2 + n_3 + n_4 = q(A) - n_1$ and $n_1 \geq 2$, by (14),

$$\begin{aligned} n = 6\ell + 4 &= n_1 + 2(n_2 + n_3 + n_4) + n_3 + 2n_4 \\ &= n_1 + 2(q(A) - n_1) + n_3 + 2n_4 \\ &= 2q(A) - n_1 + n_3 + 2n_4 \\ &\leq 2q(A) - 2 + n_3 + 2n_4. \end{aligned} \quad (15)$$

By (11), (13) and (15), we have

$$6\ell + 4 \leq 2q(A) - 2 + \frac{3\ell + n_4}{2} \leq 2q(A) - 2 + \frac{3\ell + 1}{2}. \quad (16)$$

By solving (16) for $q(A)$, and using $d(W) = 2\ell + 2$, we obtain

$$q(A) \geq \frac{9\ell}{4} + \frac{11}{4} = d(W) + 1 + \frac{\ell - 1}{4} = \frac{9d(W)}{8} + \frac{1}{2}.$$

Since A is an arbitrary matrix in $S(W)$, the result follows. $\blacksquare$

Next, we generalize Theorem 16 to whirls with more legs. The (k, ℓ) -*whirl* W for $k \geq 2$ and $\ell \geq 1$ is the tree on $n = 2k\ell + k + 1$ vertices with the axis vertex v of degree k and $2k$ legs $\{\alpha_1^i, \alpha_2^i\}_{i=1}^k$, each with ℓ vertices as illustrated in Figure 12. Note that $d(W) = 2\ell + 2$. If the number of vertices of each leg of W is not specified, we say that W is a k -*whirl*. If the numbers of the vertices of $2k$ legs are not necessarily equal, then W is a *generalized k -whirl*.




Figure 12.

In order to prove a generalization of the result in Theorem 16 for the (k, ℓ) -whirl, we first prove the following.

Lemma 17 *Suppose that W is a generalized k -whirl ($k \geq 2$) on n vertices with $2k$ legs $\{\alpha_1^i, \alpha_2^i\}_{i=1}^k$ such that each leg has at least one vertex. Let $A \in S(W)$, and let A' be the direct sum of $A[\alpha_j^i]$ for all $i \in \{1, \dots, k\}$, $j \in \{1, 2\}$. If n_r denotes the number of eigenvalues of A with multiplicity r , then the following hold:*

- (a) $n_{k+1} \leq 1$, and $n_j = 0$ for $j \geq k+2$;
- (b) If $\lambda \in \sigma(A)$ and $m_A(\lambda) = k+1$, then λ is a simple eigenvalue of $A[\alpha_j^i]$ for all $i \in \{1, \dots, k\}$, $j \in \{1, 2\}$, and $m_{A'}(\lambda) = 2k$;
- (c) If $\mu \in \sigma(A)$ and $m_A(\mu) = k$, then, for all $i \neq s$ and all j and t , μ is a simple eigenvalue of at least one of $A[\alpha_j^i]$, $A[\alpha_t^s]$, and $m_{A'}(\mu) \geq 2k-2$; and
- (d) $(2k-2)n_k + (2k)n_{k+1} \leq n - (k+1)$.

Proof. (a) By applying Proposition 13 to the $k+1$ paths illustrated in Figure 13, we conclude $p(W) = k+1$.




Figure 13.

Hence, Theorem 11 implies that there is no eigenvalue of A with multiplicity $k+2$ or more, that is, $n_j = 0$ for $j \geq k+2$. Furthermore, since the k disjoint paths $P_1, \dots, P_k$ in W in Figure 13 cover all of the vertices of W except v , Corollary 8 implies that there exists at most one eigenvalue of A with multiplicity $k+1$, that is, $n_{k+1} \leq 1$. This proves (a).

(b) Suppose $\lambda \in \sigma(A)$ and $m_A(\lambda) = k+1$. Consider the following k disjoint paths in W : $P_1 = \alpha_2^1(v_1 - v - v_k)\alpha_2^k$, $P_i = \alpha_1^i v_i \alpha_2^i$ for $i = 2, \dots, k-1$, and $P_k = \alpha_1^k$. Then $W \setminus (P_1 \cup \dots \cup P_k)$ is α_1^1 and hence, Corollary 7 implies $\lambda \in A[\alpha_1^1]$. Since α_1^1 is a path, by Theorem 11, λ is a simple eigenvalue of $A[\alpha_1^1]$. Similarly, λ is a simple eigenvalue of $A[\alpha_2^1]$ and $A[\alpha_j^i]$ for each $i = 2, \dots, k$ and $j = 1, 2$. Therefore, $m_{A'}(\lambda) = 2k$, and (b) holds.

(c) Suppose $\mu \in \sigma(A)$ and $m_A(\mu) = k$. Let $P_1 = \alpha_2^1(v_1 - v - v_k)\alpha_2^k$, and $P_i = \alpha_1^i v_i \alpha_2^i$ for $i = 2, \dots, k-1$. Then $A[W \setminus (P_1 \cup \dots \cup P_{k-1})]$ is, up to permutation similarity, equal to $A[\alpha_1^1] \oplus A[\alpha_1^k]$. Thus, by Corollary 7, μ is an eigenvalue of $A[\alpha_1^1] \oplus A[\alpha_1^k]$ and hence, by Theorem 11, μ is a simple eigenvalue of at least one of $A[\alpha_1^1]$, $A[\alpha_1^k]$. Similarly, μ is a simple eigenvalue of at least one of $A[\alpha_j^i]$, $A[\alpha_t^s]$ for all $i, s \in \{1, \dots, k\}$, $i \neq s$ and $j, t \in \{1, 2\}$. If μ is a simple eigenvalue of at most $2k-3$ of the $2k$ matrices in $\{A[\alpha_1^i], A[\alpha_2^i]\}_{i=1}^k$, then there is a pair of $A[\alpha_j^i]$, $A[\alpha_t^s]$ for $i \neq s$ such that none of $A[\alpha_j^i]$, $A[\alpha_t^s]$ has μ as an eigenvalue. Thus, μ is a simple eigenvalue of at least $2k-2$ matrices in $\{A[\alpha_1^i], A[\alpha_2^i]\}_{i=1}^k$. Therefore, $m_{A'}(\mu) \geq 2k-2$, and (c) holds.

(d) The order of A' is $n - (k+1)$. Since the number of eigenvalues of A' cannot exceed the order of A' , $(2k-2)n_k + (2k)n_{k+1} \leq n - (k+1)$. ■

Theorem 18 *Let W be the (k, ℓ) -whirl with $k \geq 3$ and $\ell \geq 2$. Then*

$$q(W) \geq d(W) + 1 + \frac{(k-2)(\ell-1)}{(k-1)^2} > d(W) + 1.$$

Proof. Let $A \in S(W)$, and let n_j be the number of eigenvalues of A with multiplicity j . By Lemma 17 (a),

$$n = 2k\ell + k + 1 = \sum_{i=1}^{k+1} i \cdot n_i.$$

Moreover,

$$\begin{aligned}
2k\ell + k + 1 &= \sum_{i=1}^{k+1} i \cdot n_i \\
&= n_1 + 2n_2 + \cdots + kn_k + (k+1)n_{k+1} \\
&\leq 2 + (k-1)(n_1 - 2 + n_2 + \cdots + n_k + n_{k+1}) + n_k + 2n_{k+1} \\
&= 2 + (k-1)(q(A) - 2) + n_k + 2n_{k+1}.
\end{aligned} \tag{17}$$

Next, since $n = 2k\ell + k + 1$, Lemma 17 (d) implies

$$(2k-2)n_k + (2k)n_{k+1} \leq 2k\ell. \tag{18}$$

Thus, by solving (18) for n_k , we have

$$n_k \leq \frac{k\ell - kn_{k+1}}{k-1}.$$

Furthermore, by (17),

$$\begin{aligned}
2k\ell + k + 1 &\leq 2 + (k-1)(q(A) - 2) + \frac{k\ell - kn_{k+1}}{k-1} + 2n_{k+1} \\
&= (k-1)q(A) - 2k + 4 + \frac{k\ell + (k-2)n_{k+1}}{k-1}.
\end{aligned}$$

By Lemma 17 (a), $n_{k+1} \leq 1$ and hence,

$$2k\ell + 3k - 3 \leq (k-1)q(A) + \frac{k\ell + k - 2}{k-1}. \tag{19}$$

By solving (19) for $q(A)$, we have

$$\begin{aligned}
q(A) &\geq \frac{[(2k\ell + 3k - 3)(k-1)] - (k\ell + k - 2)}{(k-1)^2} \\
&= \frac{2k^2\ell + 3k^2 - 3k\ell - 7k + 5}{(k-1)^2} \\
&= 2\ell + 3 + \frac{(k-2)(\ell-1)}{(k-1)^2}.
\end{aligned}$$

Since $d(W) = 2\ell + 2$, we have

$$q(A) \geq d(W) + 1 + \frac{(k-2)(\ell-1)}{(k-1)^2}. \tag{20}$$

Since A is an arbitrary matrix in $S(W)$, (20) implies

$$q(W) \geq d(W) + 1 + \frac{(k-2)(\ell-1)}{(k-1)^2} > d(W) + 1$$

for $k \geq 3$ and $\ell \geq 2$. ■

6 $q(G)$ for a Class of Connected Graphs G

In this section we apply the Smith Normal Form approach to a class of connected graphs G , and find a lower bound on $q(G)$.

Let G be a connected graph on n vertices with 6 legs L_i , each with ℓ vertices for $\ell \geq 1$, (see Figure 14) such that H is a connected graph on m vertices containing v_1, v_2, v_3 , and for all s, t , and k with $\{s, t, k\} = \{1, 2, 3\}$ there exists a unique shortest path from v_s to v_t which does not pass v_k . We use i_u, j_v to denote the pendant vertices of the 6 legs (see Figure 14).




Figure 14.

Using the Smith Normal Form approach, we show the following.

Theorem 19 *Let G be the connected graph described in Figure 14. Then*

$$q(G) \geq \frac{9\ell}{4} - 2m + \frac{15}{2}.$$

Proof. Let $A \in S(G)$, and $\lambda_1, \dots, \lambda_q$ be the distinct eigenvalues of A , and let n_j be the number of eigenvalues of A with multiplicity j . We consider $xI - A$. Note that $D(xI - A)$ has a loop at each vertex.

Now we compute the determinant of an $(n - 3)$ by $(n - 3)$ submatrix of $xI - A$, and find an upper bound on $\sum_{\lambda_j: m_A(\lambda_j) \geq 4} (m_A(\lambda_j) - 2)$. Let M be the matrix obtained from

$xI - A$ by replacing column i_s by e_{j_s} for each $s = 1, 2, 3$. In terms of digraphs, $D(M)$ is obtained from $D(xI - A)$ by deleting all incoming arcs to vertices i_1, i_2, i_3 , and inserting the arcs $(j_1, i_1), (j_2, i_2), (j_3, i_3)$, each with weight 1. Hence, (j_s, i_s) is the unique arc of $D(M)$ ending at vertex i_s for each $s = 1, 2, 3$. Moreover, if there exists a directed walk from i_s to j_s in $D(M)$ which is not the directed path $P_s = (i_s, \dots, v_s, \dots, j_s)$, then the directed walk has a loop or it repeats v_s at least twice. Thus, $\beta_s = (P_s, (j_s, i_s))$ is the unique directed cycle containing (j_s, i_s) for each $s = 1, 2, 3$. Therefore, each set of disjoint directed cycles of $D(M)$ that cover every vertex consists of the directed cycles β_s 's along with disjoint directed cycles covering every vertex of $H \setminus U$ where $U = \{v_1, v_2, v_3\}$. By (5),

this implies that

$$\det M = \pm \prod_{s=1}^3 \text{wt}(P_s) \cdot \det M[H \setminus U]. \quad (21)$$

Note that $M[H \setminus U] = (xI - A)[H \setminus U]$, and $\text{wt}(P_s)$ is a nonzero constant for each $s = 1, 2, 3$. By Laplace expansion of the determinant along the columns i_1, i_2, i_3 of M , $\det M = \pm \det[(xI - A)(\{j_1, j_2, j_3\}, \{i_1, i_2, i_3\})]$. Thus, by (21),

$$\det[(xI - A)(\{j_1, j_2, j_3\}, \{i_1, i_2, i_3\})] = c \cdot \det((xI - A)[H \setminus U])$$

for some nonzero constant c .

Since $(xI - A)(\{j_1, j_2, j_3\}, \{i_1, i_2, i_3\})$ is an $n - 3$ by $n - 3$ submatrix of $xI - A$, we have

$$\Delta_{n-3}(x) | \det((xI - A)[H \setminus U]).$$

Thus, by Theorem 3, if λ is an eigenvalue of A with $m_A(\lambda) \geq 4$, then λ is a zero of $(xI - A)[H \setminus U]$ with multiplicity $m_A(\lambda) - 3$ or more. Since the order of $(xI - A)[H \setminus U]$ is $m - 3$, we have

$$\sum_{\lambda_j: m_A(\lambda_j) \geq 4} 1 \leq \sum_{\lambda_j: m_A(\lambda_j) \geq 4} (m_A(\lambda_j) - 3) \leq m - 3. \quad (22)$$

Thus, by (22),

$$\sum_{\lambda_j: m_A(\lambda_j) \geq 4} (m_A(\lambda_j) - 2) = \sum_{\lambda_j: m_A(\lambda_j) \geq 4} (m_A(\lambda_j) - 3) + \sum_{\lambda_j: m_A(\lambda_j) \geq 4} 1 \leq 2m - 6. \quad (23)$$

Next, we compute the determinants of $(n - 2)$ by $(n - 2)$ submatrices of $xI - A$ and thereby, find an upper bound on n_3 . Let N be the matrix obtained from $xI - A$ by replacing column a_s by e_{b_t} , and column i_k by e_{j_k} where $a, b \in \{i, j\}$ and $\{s, t, k\} = \{1, 2, 3\}$. In terms of digraphs, $D(N)$ is obtained from $D(xI - A)$ by deleting all incoming arcs to a_s, i_k , and inserting $(b_t, a_s), (j_k, i_k)$, each with weight 1 (see Figure 15).




Figure 15.

In Figure 15, edges represent directed 2-cycles. Therefore, $D(N)$ has exactly one arc, namely (b_t, a_s) , ending at a_s and exactly one arc, namely (j_k, i_k) , ending at i_k . Moreover, each directed path from a_s to b_t passes through v_s and v_t . Let $Q_1^k, \dots, Q_{d_k}^k$ be the directed paths from v_s to v_t not containing vertex v_k , and Q_1^k be the unique shortest directed path from v_s to v_t that does not go through v_k . Then, for each $g = 1, \dots, d_k$, $(P_{a_s \rightarrow v_s}, Q_g^k, P_{v_t \rightarrow b_t})$ is a directed path from a_s to b_t not containing vertex v_k , and $(P_{a_s \rightarrow v_s}, Q_1^k, P_{v_t \rightarrow b_t})$ is the unique shortest directed path from a_s to b_t that does not go through vertex v_k . Note that there exists exactly one directed cycle α of $D(N)$ containing (j_k, i_k) , $\alpha = (P_{i_k \rightarrow v_k}, P_{v_k \rightarrow j_k}, (j_k, i_k))$. Thus, the directed cycles $\beta_g^k = (P_{a_s \rightarrow v_s}, Q_g^k, P_{v_t \rightarrow b_t}, (b_t, a_s))$ ($g = 1, \dots, d_k$) are the only directed cycles containing (b_t, a_s) , which are disjoint from α . Let $H' = H \setminus \{v_k\}$. Then, by (5),

$$\det N = \text{swt}(\alpha) \cdot \det N[L] \cdot \det N[R] \cdot \sum_{g=1}^{d_k} \text{swt}(\beta_g^k) \cdot \det N[H' \setminus \beta_g^k], \quad (24)$$

where $N[H' \setminus \beta_g^k]$ is the principal submatrix of N whose rows and columns correspond to $V(H') \setminus V(\beta_g^k)$.

Since $H' \setminus \beta_1^k$ is the unique one among $(H' \setminus \beta_g^k)$'s having the largest number of vertices and $\text{swt}(\beta_g^k)$'s are nonzero constants, $\sum_{g=1}^{d_k} \text{swt}(\beta_g^k) \cdot \det N[H' \setminus \beta_g^k]$ is a nonzero polynomial of degree $|V(H') \setminus V(\beta_1^k)|$.

Note that $\text{swt}(\beta_g^k) = (-1)^{v(P_{a_s \rightarrow v_s}) + v(Q_g^k) + v(P_{v_t \rightarrow b_t}) - 3} \text{wt}(P_{a_s \rightarrow v_s}) \cdot \text{wt}(Q_g^k) \cdot \text{wt}(P_{v_t \rightarrow b_t})$, and $V(H') \setminus V(\beta_g^k) = V(H') \setminus V(Q_g^k)$, where $v(P_{a_s \rightarrow v_s})$, $v(Q_g^k)$ and $v(P_{v_t \rightarrow b_t})$ are the numbers of the vertices of those directed paths. Hence, from (24),

$$\begin{aligned} \det N &= \text{swt}(\alpha) \cdot \det N[L] \cdot \det N[R] \cdot (-1)^{v(P_{a_s \rightarrow v_s}) + v(P_{v_t \rightarrow b_t}) - 3} \text{wt}(P_{a_s \rightarrow v_s}) \cdot \text{wt}(P_{v_t \rightarrow b_t}) \\ &\quad \cdot \sum_{g=1}^{d_k} (-1)^{v(Q_g^k)} \text{wt}(Q_g^k) \cdot \det N[H' \setminus Q_g^k]. \end{aligned}$$

Since $\text{swt}(\alpha)$, $\text{wt}(P_{a_s \rightarrow v_s})$ and $\text{wt}(P_{v_t \rightarrow b_t})$ are nonzero constants,

$$\det N = c \cdot \det N[L] \cdot \det N[R] \cdot \sum_{g=1}^{d_k} (-1)^{v(Q_g^k)} \text{wt}(Q_g^k) \cdot \det N[H' \setminus Q_g^k], \quad (25)$$

for some nonzero constant c .

Note that $\det N[L] = \det((xI - A)[L])$, and $\det N[R] = \det((xI - A)[R])$. By Laplace expansion of the determinant along the columns a_s, i_k of N , $\det N = \pm \det[(xI - A)(\{b_t, j_k\}, \{a_s, i_k\})]$. Thus, by (25),

$$\det[(xI - A)(\{b_t, j_k\}, \{a_s, i_k\})] = \pm c \cdot \det N[L] \cdot \det N[R] \cdot f_k(x)$$

where $f_k(x) = \sum_{g=1}^{d_k} (-1)^{v(Q_g^k)} \text{wt}(Q_g^k) \cdot \det N[H' \setminus Q_g^k]$. Note that $f_k(x)$ is independent of the choices of a_s, b_t . Since $(xI - A)(\{b_t, j_k\}, \{a_s, i_k\})$ is an $n-2$ by $n-2$ submatrix of $xI - A$,

$$\Delta_{n-2}(x) | \det N[L] \cdot \det N[R] \cdot f_k(x).$$

Let $\mu \in \sigma(A)$ and $m_A(\mu) = 3$. Then, by Theorem 3, μ is a zero of $\det N[L] \cdot \det N[R] \cdot f_k(x)$. If μ is an eigenvalue of neither $A[L]$ nor $A[R]$, then μ is a zero of $f_k(x)$. Note that L and R are arbitrary legs of G incident to different v_i 's.

If μ is not an eigenvalue of at least 4 of $A[L_r]$'s, then there exist $A[L_u]$ and $A[L_v]$ such that $\lambda \notin \sigma(A[L_u])$ and $\lambda \notin \sigma(A[L_v])$, and L_u, L_v are connected to different v_i 's. Thus, in this case, μ is a zero of $f_k(x)$ for some $k \in \{1, 2, 3\}$. Note that $V(H') \setminus V(Q_g^k) \subseteq V(H \setminus U)$ and thereby, the degree of $f_k(x)$ is $m-3$ or less for each $k = 1, 2, 3$. Therefore, the number, n'_3 , of such μ satisfies

$$n'_3 \leq 3(m-3). \quad (26)$$

Let n''_3 be the number of eigenvalues μ of A with multiplicity 3 such that μ is an eigenvalue of at least 4 of $A[L_r]$'s. Then $n_3 = n'_3 + n''_3$. Since μ is an eigenvalue of $A[L_1] \oplus \cdots \oplus A[L_6]$ with multiplicity at least 4, and the order of $A[L_1] \oplus \cdots \oplus A[L_6]$ is 6ℓ , $4n''_3 \leq 6\ell$. Equivalently,

$$n''_3 \leq \frac{3\ell}{2}. \quad (27)$$

Now, we compute a lower bound on $q(A) = q$. Note that

$$\begin{aligned} n = 6\ell + m = \sum_{i=1}^q m_A(\lambda_i) &= 2q - n_1 + n'_3 + n''_3 + \sum_{\lambda_j: m_A(\lambda_j) \geq 4} (m_A(\lambda_j) - 2) \\ &\leq 2q + n'_3 + n''_3 + \sum_{\lambda_j: m_A(\lambda_j) \geq 4} (m_A(\lambda_j) - 2). \end{aligned}$$

By (23), (26), and (27), we have

$$6\ell + m \leq 2q + 3(m-3) + \frac{3\ell}{2} + 2m - 6. \quad (28)$$

Hence, by solving (28) for $q = q(A)$, we have

$$q(A) \geq \frac{9\ell}{4} - 2m + \frac{15}{2}. \quad (29)$$

Since A is an arbitrary matrix in $S(G)$, (29) implies

$$q(G) \geq \frac{9\ell}{4} - 2m + \frac{15}{2}. \quad \blacksquare$$

References

- [1] F. Barioli and S.M. Fallat, On two conjectures regarding an inverse eigenvalue problem for acyclic symmetric matrices, *Electronic J. Lin. Alg.* 11 (2004), 41-50.
- [2] W.G. Bridges and R. Mena, Lecture Notes, University of Wyoming, 1984.
- [3] R.A. Brualdi, and H.J. Ryser, *Combinatorial Matrix Theory*, Cambridge University Press, Cambridge (1991).
- [4] M. Fiedler, Eigenvectors of acyclic matrices, *Czech. Math. J.* 25 (1975), 607-618.
- [5] C. Godsil and G. Royle, *Algebraic Graph Theory*, Springer (2001).
- [6] B. Hartley and T.O. Hawkes, *Rings, Modules and Linear Algebra*, Chapman & Hall (1994).
- [7] R. Horn and C.R. Johnson, *Matrix Analysis*, Cambridge University Press, New York (1985).
- [8] C.R. Johnson and A. Leal Duarte, The maximum multiplicity of an eigenvalue in a matrix whose graph is a tree, *Lin. Mult. Alg.* 46 (1999), 139-144.

- [9] C.R. Johnson and A. Leal Duarte, On the possible multiplicities of the eigenvalues of a Hermitian matrix whose graph is a tree, *Lin. Alg. and Appl.* 348 (2002), 7-21.
- [10] C.R. Johnson, A. Leal Duarte and C.M. Saiago, The Parter-Wiener theorem: Refinement and Generalization, *SIAM J. Matrix Anal. Appl.* 25 (2003), 352-361.
- [11] C.R. Johnson, A. Leal Duarte and C.M. Saiago, Inverse eigenvalue problems and list of eigenvalues for matrices whose graph is a tree: the case of generalized stars and double generalized stars, *Lin. Alg. and Appl.* 373 (2003), 311-330.
- [12] C.R. Johnson, A. Leal Duarte, C.M. Saiago, B.D. Sutton, and A.J. Witt, On the relative position of multiple eigenvalues in the spectrum of an Hermitian matrix with a given graph, *Lin. Alg. and Appl.* 363 (2003), 147-159.
- [13] A. Leal Duarte, Construction of acyclic matrices from spectral data, *Lin. Alg. and Appl.* 113 (1989), 173-182.
- [14] A. Leal Duarte and C.R. Johnson, On the minimum number of distinct eigenvalues for a symmetric matrix whose graph is a given tree, *Math. Inequal. Appl.* 5 (2002), 175-180.
- [15] P.M. Nylén, Minimum-Rank Matrices With Prescribed Graph, *Lin. Alg. and Appl.* 248 (1996), 303-316.